arXiv:1511.00230v1 [math.FA] 1 Nov 2015

Volterra-composition operators on the weighted Bergman space with exponential type weights

Pan Xiaohua[1], Wu Yan[2], Xu Xianmin[2]

1. Department of Mathematics, Zhejiang Normal University,
Jinhua, Zhejiang 321004, P. R. China

2. College of Mathematics Physics and Information Engineering,
Jiaxing University, Jiaxing, Zhejiang 314001, P. R. China

Abstract. The properties of Volterra-composition operators on the weighted Bergman space with exponential type weights are investigated in this paper. For a certain class of subharmonic function $\psi : \mathbb{D} \to \mathbb{R}$, we state some necessary and sufficient conditions that a Volterra-composition operator V_φ^g from the weighted Bergman space $AL^2_\psi(\mathbb{D})$ to Bloch type space $B_\psi(\mathbb{D})$ (or little Bloch type space $B_{\psi,0}(\mathbb{D})$) must satisfy for V_φ^g to be bounded or compact.

Keywords The weighted Bergman space; Bloch type space(little Bloch type space); Volterra-composition operator; bounded; compact.

MR(2010) Subject Classification 74D05; 35Q72; 35L70; 35B40

[1]

1 Introduction

Let $\mathbb{D} = \{z : |z| < 1\}$ be the open unit disk in the complex plane $\mathbb{C}$ and dA be the normalized area measure on $\mathbb{D}$. Let $L^2(\mathbb{D}, dA)$ be the space of square integrable functions and let $L^\infty(\mathbb{D}, dA)$ be the space of essential bounded measurable functions. We will use abbreviations $L^2(\mathbb{D})$ for $L^2(\mathbb{D}, dA)$ and $L^\infty(\mathbb{D})$ for $L^\infty(\mathbb{D}, dA)$.

[1] This research was supported by the National Natural Science Foundation of China under Grant (No.11301224,11401256,11501249)

For a subharmonic function $\psi : \mathbb{D} \to \mathbb{R}$, let $L^\infty_\psi(\mathbb{D})$ be the space of measurable functions f on $\mathbb{D}$ such that $e^{-\psi} f \in L^\infty(\mathbb{D})$ and let $L^2_\psi(\mathbb{D})$ be the Hilbert space of measurable function f on $\mathbb{D}$ such that

$$\|f\|_{L^2_\psi} = (\int_{\mathbb{D}} |f|^2 e^{-2\psi} dA)^{\frac{1}{2}} < \infty$$

The inner product of $L^2_\psi(\mathbb{D})$ is given by

$$\langle f, g \rangle_{L^2_\psi} = \int_{\mathbb{D}} f\overline{g} e^{-2\psi} dA$$

Let $H^\infty_\psi(\mathbb{D})$ be the subspace of $L^\infty_\psi(\mathbb{D})$ consisting of analytic functions and $AL^2_\psi(\mathbb{D})$ be the closed subspace of $L^2_\psi(\mathbb{D})$ consisting of analytic functions. $AL^2_\psi(\mathbb{D})$ is called the weighted Bergman space with exponential type weights(see [3],[4]).

An analytic function f is said to belong to the Bloch-type space $B_\psi(\mathbb{D})$ if

$$b_\psi(f) = \sup_{z\in\mathbb{D}} e^{-2\psi(z)} |f'(z)| < \infty.$$

Let $B_{\psi,0}(\mathbb{D})$ denote the subspace of $B_\psi(\mathbb{D})$ such that

$$\lim_{|z|\to 1} e^{-2\psi(z)} |f'(z)| = 0.$$

This space is called little Bloch-type space.

Definition 1.1. [1],[2] For real valued function $\psi \in C^2(\mathbb{D})$ with $\Delta\psi > 0$, where Δ is the Laplace operator. Let $\tau(z) = (\Delta\psi(z))^{-\frac{1}{2}}$. We say that $\psi \in \mathcal{D}$ if the following conditions are satisfied.
(1) There exists a constant $C_1 > 0$ such that $|\tau(z) - \tau(\xi)| \le C_1|z - \xi|$ for $z, \xi \in \mathbb{D}$.
(2) There exists a constant $C_2 > 0$ such that $\tau(z) \le C_2(1 - |z|)$ for $z \in \mathbb{D}$.
(3) There exist constants $0 < C_3 < 1$ and $a > 0$ such that $\tau(z) \le \tau(\xi) + C_3(1 - |z|)$ for $|z - \xi| > a\tau(\xi)$.

Let $H(\mathbb{D})$ be the space of analytic functions on $\mathbb{D}$ and suppose that $g : \mathbb{D} \to \mathbb{C}$ is an analytic function, φ is an analytic self-map of the unit disk and $f \in H(\mathbb{D})$, the Riemann-Stieltjes operator is defined by

$$T_g f(z) = \int_0^1 f(tz) z g'(tz) dt = \int_0^z f(t) g'(t) dt, \qquad z \in \mathbb{D}.$$

The composition operator C_φ is defined as $C_\varphi f \triangleq f \circ \varphi$.

Let V^g_φ be the Volterra-composition operator which is defined as

$$V^g_\varphi f(z) \triangleq (T_g \circ C_\varphi f)(z) = \int_0^z f(\varphi(t)) g'(t) dt, \quad \forall f \in H(\mathbb{D}),\ z \in \mathbb{D}.$$

Note that when φ is an analytic self-map of the unit disk and $f, g \in H(\mathbb{D})$, we have

$$(V_\varphi^g f)^{'}(z) = f(\varphi(z))g^{'}(z).$$

In this paper, we will characterize the boundedness and compactness of the operator V_φ^g from the weighted Bergman space with exponential type weights to the Bloch-type space (or little Bloch-type space). These properties of Hankel operators have been considered(see [1],[2]).

We assume that $H_\psi^\infty(\mathbb{D})$ is dense in $AL_\psi^2(\mathbb{D})$, the set of all polynomials is dense in $AL_\psi^2(\mathbb{D})$.

2 Preliminaries

In order to prove the main results, we need the following lemmas.

Lemma 2.1. *Let*

$$\|f\|_{B_\psi} = |f(0)| + \sup_{z\in\mathbb{D}} e^{-2\psi(z)}|f^{'}(z)|,$$

then Bloch–type space $B_\psi(\mathbb{D})$ is a Banach space with the norm $\|\cdot\|_{B_\psi}$.

Proof. At first, we will show that $\|\cdot\|_{B_\psi}$ is a norm. For a function $f \in B_\psi(\mathbb{D})$, we define

$$b_\psi(f) = \sup_{z\in\mathbb{D}} e^{-2\psi(z)}|f^{'}(z)|.$$

It is easy to see that

$$\|\alpha f\|_{B_\psi} = |\alpha f(0)| + b_\psi(\alpha f) = |\alpha||f(0)| + |\alpha| b_\psi(f) = |\alpha|\|f\|_{B_\psi}.$$

$$\begin{aligned} \|f+g\|_{B_\psi} &= |f(0)+g(0)| + b_\psi(f+g) \le |f(0)| + b_\psi(f) + |g(0)| + b_\psi(g) \\ &= \|f\|_{B_\psi} + \|g\|_{B_\psi}. \end{aligned}$$

So $\|\cdot\|_{B_\psi}$ is a semi-norm.

Assume that $\|f\|_{B_\psi} = |f(0)| + b_\psi(f) = 0$, we have $f(0) = 0$ and $b_\psi(f) = 0$, which means that

$$\sup_{z\in\mathbb{D}} e^{-2\psi(z)}|f^{'}(z)| = 0.$$

Since $e^{-2\psi(z)} \neq 0$ when $z \in \mathbb{D}$, it must be $|f'(z)| \equiv 0$. It follows that $f(z) \equiv C$ for some constant C. Since $f(0) = 0$, we obtain $C = 0$. It implies that $f = 0$.

Now we are going to prove the completeness of $\|\cdot\|_{B_\psi}$. Suppose that $\{f_n\}_{n\in\mathbb{N}}$ is a Cauchy sequence of $B_\psi(\mathbb{D})$, i.e.

$$\|f_n - f_m\|_{B_\psi} = |f_n(0) - f_m(0)| + \sup_{z\in\mathbb{D}} e^{-2\psi(z)}|f_n^{'}(z) - f_m^{'}(z)| \to 0 \quad (n, m \to \infty).$$

Since

$$\begin{aligned} |f_n(z) - f_m(z)| &= |\int_0^z [f_n^{'}(t) - f_m^{'}(t)]dt + [f_n(0) - f_m(0)]| \\ &\leq \int_0^z |f_n^{'}(t) - f_m^{'}(t)|dt + |f_n(0) - f_m(0)|, \end{aligned}$$

for each $z \in \mathbb{D}$, there exists a $f \in H(\mathbb{D})$ such that f_n uniformly converges to f on any compact subsets of $\mathbb{D}$. By Cauchy integral formula, we have $f_n^{'}$ uniformly converges to $f^{'}$ on any compact subsets of $\mathbb{D}$.

Since $b_\psi(f_n - f_m) \to 0$, for any $\varepsilon > 0$, there exists a positive integer N, such that when $n, m > N$, we have

$$e^{-2\psi(z)}|f_n^{'}(z) - f_m^{'}(z)| < \varepsilon, \quad for\ all\ z \in \mathbb{D}.$$

Letting $m \to \infty$ in the above inequality, we can see that

$$e^{-2\psi(z)}|f_n^{'}(z) - f^{'}(z)| \leq \varepsilon, \quad for\ all\ z \in \mathbb{D},$$

this implies that $b_\psi(f_n - f) \to 0$ $(n \to \infty)$. It follows that $\|f_n - f\|_{B_\psi} \to 0$ $(n \to \infty)$. It is easy to see that there exists a positive integer n such that for all $z \in \mathbb{D}$

$$e^{-2\psi(z)}|f^{'}(z)| \leq 1 + e^{-2\psi(z)}|f_n^{'}(z)| \leq 1 + b_\psi(f_n) < \infty.$$

i.e. $f \in B_\psi$, thus the completeness is proved.

Therefore, $B_\psi(\mathbb{D})$ is a Banach space with the norm $\|\cdot\|_{B_\psi}$. □

Notation: we write $f(z, w) \sim g(z, w)$ if there exist positive constants C and C' such that

$$Cg(z, w) \leq f(z, w) \leq C'g(z, w)$$

.

Lemma 2.2. [2] *Let $\psi \in \mathcal{D}$, then we have*

$$K(z, z)e^{-2\psi(z)} \sim (\tau(z))^{-2} = \Delta\psi(z), \quad z \in \mathbb{D}.$$

where $K(z, w)$ is the reproducing kernel of $AL^2_\psi(\mathbb{D})$.

Lemma 2.3. *Let $f \in AL^2_\psi(\mathbb{D})$ and $\psi \in \mathcal{D}$, then*

$$|f(z)| \leq \sqrt{K(z, z)}\|f\|_{L^2_\psi}.$$

Proof. $|f(z)| = |\langle f, K_z\rangle| \leq \|f\|_{L^2_\psi}\|K_z\|_{L^2_\psi} = \sqrt{K(z, z)}\|f\|_{L^2_\psi}$. □

Lemma 2.4. *Let φ be an analytic self-map of the unit disk $\mathbb{D}$ and $g \in H(\mathbb{D})$. Assume that $V_\varphi^g : AL_\psi^2(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded. Then V_φ^g is compact if and only if $V_\varphi^g f_k$ converges to zero as $k \to \infty$, for any bounded sequence $\{f_k\}_{k\in\mathbb{N}}$ in $AL_\psi^2(\mathbb{D})$ which uniformly converges to zero on compact subsets of $\mathbb{D}$.*

Proof. First, assume that V_φ^g is compact. Let $\{f_k\}_{k\in\mathbb{N}}$ be any bounded sequence in $AL_\psi^2(\mathbb{D})$ and uniformly converges to zero on compact subsets of $\mathbb{D}$. Since V_φ^g is a compact operator , there exists a subsequence of $\{f_k\}_{k\in\mathbb{N}}$ (without loss of generality, we assume it is $\{f_k\}_{k\in\mathbb{N}}$) and a function $h \in B_\psi(\mathbb{D})$ such that

$$\|V_\varphi^g f_k - h\|_{B_\psi} \to 0 (k \to \infty).$$

That is,

$$|V_\varphi^g f_k(0) - h(0)| + \sup_{z\in\mathbb{D}} e^{-2\psi(z)} |f_k(\varphi(z))g^{'}(z) - h^{'}(z)| \to 0 (k \to \infty).$$

By the definition of V_φ^g, it is obvious that $V_\varphi^g f_k(0) = 0$. It follows that $h(0) = 0$. It is not difficult to see that $f_k(\varphi(z))g^{'}(z)$ uniformly converges to $h^{'}(z)$ on any compact subsets of $\mathbb{D}$. Besides, $\{f_k\}_{k\in\mathbb{N}}$ uniformly converges to zero on any compact subsets of $\mathbb{D}$ and $g \in H(\mathbb{D})$, it follows that $h^{'}(z) \equiv 0$. Also, as $h(0) = 0$, we have $h(z) \equiv 0$, that is

$$\|V_\varphi^g f_k\|_{B_\psi} \to 0 (k \to \infty).$$

Conversely, let $\{f_k\}_{k\in\mathbb{N}}$ be any bounded sequence in $AL_\psi^2(\mathbb{D})$. By Lemma 2.3, we have $|f_k(z)| \le \sqrt{K(z,z)}\|f_k\|_{L_\psi^2}$. therefore, $\{f_k\}_{k\in\mathbb{N}}$ is uniformly bounded on any compact subset K of $\mathbb{D}$.

By Montel Theorem, there exists a subsequence $\{f_{k_n}\}_{k_n\in\mathbb{N}}$ of $\{f_k\}_{k\in\mathbb{N}}$ and an analytic function f, such that $\{f_{k_n}\}$ converges to f uniformly on any compact subsets K of the unit disk $\mathbb{D}$.

According to the Fatou Lemma, $f \in AL_\psi^2(\mathbb{D})$. Together with the assumption, we have $\|V_\varphi^g(f_{k_n} - f)\|_{B_\psi} \to 0$ $(n \to \infty)$. Therefore, $\{V_\varphi^g f_k\}_{k\in\mathbb{N}}$ has subsequence $\{V_\varphi^g f_{k_n}\}_{n\in\mathbb{N}}$ which converges to $V_\varphi^g f \in AL_\psi^2(\mathbb{D})$. From the characterization of compact operators, we see that $V_\varphi^g : AL_\psi^2(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact. □

Lemma 2.5. *Let $\psi \in \mathcal{D}$, then the bounded closed set K of $B_{\psi,0}(\mathbb{D})$ is compact if and only if $\limsup_{|z|\to 1 f\in K} e^{-2\psi(z)}|f^{'}(z)| = 0$.*

Proof. Assume that the bounded closed set K of $B_{\psi,0}(\mathbb{D})$ is compact. Let $\varepsilon > 0$, we have $\bigcup_{f\in K} B(f, \frac{\varepsilon}{2}) \supset K$, where $B(f, \frac{\varepsilon}{2})$ is a ball with center f and radius $\frac{\varepsilon}{2}$. Because K is compact, there exist $f_1, f_2, \cdots, f_n \in K$, such that for any $f \in K$, we have

$$\|f - f_j\|_{B_\psi} < \frac{\varepsilon}{2} \quad (1 \le j \le n, j \in \mathbb{N}^*).$$

Hence, for any $z \in \mathbb{D}$, we have, for $1 \leq j \leq n$,

$$e^{-2\psi(z)}|f^{'}(z)| \leq e^{-2\psi(z)}|f_j^{'}(z)| + \frac{\varepsilon}{2}.$$

Since $f_j \in B_{\psi,0}(\mathbb{D})$, for each positive integer j, there exists a positive real number $r_j \in (0,1)$ such that $e^{-2\psi(z)}|f_j^{'}(z)| < \frac{\varepsilon}{2}$ when $r_j < |z| < 1$. Let $r = \max\{r_1, r_2, \cdots, r_n\}$, then we have, when $r < |z| < 1$,

$$e^{-2\psi(z)}|f^{'}(z)| \leq \varepsilon.$$

Namely,

$$\limsup_{|z|\to 1 f\in K} e^{-2\psi(z)}|f^{'}(z)| = 0.$$

Conversely, suppose that the sequence $\{f_n\}_{n\in\mathbb{N}} \subset K$. Since K is bounded, there is a positive constant M such that for any $n \in \mathbb{Z}^+$, $\|f_n\|_{B_\psi} \leq M$. By Lemma 2.3, it is easy to see $\{f_n\}_{n\in\mathbb{N}}$ is uniformly bounded on any compact subsets of $\mathbb{D}$. By Montel Theorem, there exists subsequence of $\{f_n\}_{n\in\mathbb{N}}$ (without loss of generality , we suppose it to be $\{f_n\}_{n\in\mathbb{N}}$) and an analytic function f, such that $\{f_n\}_{n\in\mathbb{N}}$ converges to f uniformly on any compact subsets of $\mathbb{D}$.

Since

$$\limsup_{|z|\to 1 f\in K} e^{-2\psi(z)}|f^{'}(z)| = 0$$

and $\{f_n\}_{n\in\mathbb{N}} \subseteq K$, for any $\varepsilon > 0$, there exists an $r \in (0,1)$, such that $e^{-2\psi(z)}|f_n^{'}(z)| < \frac{\varepsilon}{2}$ for all n, when $r < |z| < 1$. Furthermore, $e^{-2\psi(z)}|f^{'}(z)| < \frac{\varepsilon}{2}$ when $r < |z| < 1$, therefore we have

$$e^{-2\psi(z)}|f_n^{'}(z) - f^{'}(z)| < \varepsilon$$

when $r < |z| < 1$.

Combines with Cauchy Integral Theorem, $f_n^{'}$ converges to $f^{'}$ uniformly on any compact subset of $\mathbb{D}$. Note that $e^{-2\psi(z)}$ is continuous on $r\overline{\mathbb{D}}$, hence for any $z \in r\overline{\mathbb{D}}$, we have $|e^{-2\psi(z)}| \leq C$ for some constant C. As $f_n^{'}$ converges to $f^{'}$ uniformly on $r\overline{\mathbb{D}}$, there exists an integer G, when $n \geq G$, $|f_n^{'}(z) - f^{'}(z)| < \frac{\varepsilon}{C}$ holds for any $z \in r\overline{\mathbb{D}}$.

Thus, it is easy to see, when $n \geq G$, that $e^{-2\psi(z)}|f_n^{'}(z) - f^{'}(z)| < \varepsilon$ holds for any $z \in \mathbb{D}$. This implies $\|f_n - f\|_{B_\psi} \to 0 \ \ (n \to \infty)$. Since K is closed, we have $f \in K$, that means K is compact. □

3 Main Results

Theorem 3.1. *Let φ be an analytic self-map of the unit disk $\mathbb{D}$ and $g \in H(\mathbb{D})$, $\psi \in \mathcal{D}$. Then $V_\varphi^g : AL_\psi^2(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded if and only if*

$$\sup_{z\in\mathbb{D}} e^{\psi(\varphi(z))-2\psi(z)}\sqrt{\Delta\psi(\varphi(z))}|g'(z)| < \infty.$$

Proof. By Lemma 2.2,

$$e^{\psi(\varphi(z))}\sqrt{\Delta\psi(\varphi(z))} \sim \sqrt{K(\varphi(z),\varphi(z))},$$

therefore, it suffices for us to prove that V_φ^g is bounded if and only if

$$\sup_{z\in\mathbb{D}}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| < \infty.$$

First, assume that $V_\varphi^g : AL_\psi^2(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded. For every $w \in \mathbb{D}$, let $f_w(z) = k_w(z) = \frac{K(z,w)}{\sqrt{K(w,w)}}$. It is easy to check that $f_w \in AL_\psi^2(\mathbb{D})$ and $\|f_w\|_{L_\psi^2} = 1$ for any $w \in \mathbb{D}$.

Hence, for a fixed $w \in \mathbb{D}$,

$$\begin{aligned} |f_w(\varphi(z))||g'(z)|e^{-2\psi(z)} &= |(V_\varphi^g f_w)'(z)|e^{-2\psi(z)} \\ &\le \|V_\varphi^g f_w\|_{B_\psi} \\ &\le \|V_\varphi^g\|\|f_w\|_{L_\psi^2} \\ &= \|V_\varphi^g\| < \infty \end{aligned}$$

for any $z \in \mathbb{D}$. Noticing that

$$f_w(\varphi(z)) = \frac{K(\varphi(z),w)}{\sqrt{K(w,w)}}, \quad \text{for all } z \in \mathbb{D},$$

by setting $w = \varphi(z)$, we have

$$f_{\varphi(z)}(\varphi(z)) = \frac{K(\varphi(z),\varphi(z))}{\sqrt{K(\varphi(z),\varphi(z))}} = \sqrt{K(\varphi(z),\varphi(z))}.$$

It follows that

$$\sup_{z\in\mathbb{D}}\sqrt{K(\varphi(z),\varphi(z))}|g'(z)|e^{-2\psi(z)} < \infty.$$

Conversely, assume that

$$\sup_{z\in\mathbb{D}}\sqrt{K(\varphi(z),\varphi(z))}|g'(z)|e^{-2\psi(z)} = M < \infty.$$

For any $f \in AL_\psi^2(\mathbb{D})$. By Lemma 2.3, we have

$$\begin{aligned} \|V_\varphi^g f\|_{B_\psi} &= |V_\varphi^g f(0)| + \sup_{z\in\mathbb{D}}|(V_\varphi^g f)'(z)|e^{-2\psi(z)} \\ &= \sup_{z\in\mathbb{D}}|f(\varphi(z))g'(z)|e^{-2\psi(z)} \\ &\le \sup_{z\in\mathbb{D}}\|f\|_{L_\psi^2}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| \\ &= \|f\|_{L_\psi^2}M. \end{aligned}$$

Therefore, V_φ^g is bounded. □

Theorem 3.2. *Let φ be an analytic self-map of the unit disk $\mathbb{D}$ and $g \in H(\mathbb{D})$, $\psi \in \mathcal{D}$. Then $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded if and only if $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded and*

$$\lim_{|z|\to 1} e^{-2\psi(z)}|g'(z)| = 0.$$

Proof. Assume that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded. It is clear that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded. Taking $f(z) = 1 \in AL^2_\psi(\mathbb{D})$ and $V_\varphi^g f \in B_{\psi,0}(\mathbb{D})$, then

$$\begin{aligned} 0 = \lim_{|z|\to 1} |(V_\varphi^g f)'(z)|e^{-2\psi(z)} &= \lim_{|z|\to 1} |f(\varphi(z))||g'(z)|e^{-2\psi(z)} \\ &= \lim_{|z|\to 1} |g'(z)|e^{-2\psi(z)} \end{aligned}$$

Conversely, suppose that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded and $\lim_{|z|\to 1} e^{-2\psi(z)}|g'(z)| = 0$. For each polynomial $p(z)$, the following inequality holds

$$\begin{aligned} |(V_\varphi^g p)'(z)|e^{-2\psi(z)} &= |p(\varphi(z))||g'(z)|e^{-2\psi(z)} \\ &\leq M_p|g'(z)|e^{-2\psi(z)}. \end{aligned}$$

where $M_p = \sup_{z\in\mathbb{D}}|p(z)|$. Since $M_p < \infty$ and $\lim_{|z|\to 1} |g'(z)|e^{-2\psi(z)} = 0$, then

$$\lim_{|z|\to 1} |(V_\varphi^g p)'(z)|e^{-2\psi(z)} = 0.$$

That means for each polynomial p, $V_\varphi^g p(z) \in B_{\psi,0}(\mathbb{D})$. Since the set consisting of polynomials is dense in $AL^2_\psi(\mathbb{D})$, for every $f \in AL^2_\psi(\mathbb{D})$, there is a sequence of polynomials $\{p_k\}_{k\in\mathbb{N}}$ such that

$$\|f - p_k\|_{L^2_\psi} \to 0 \quad (k \to \infty).$$

Hence,

$$\|V_\varphi^g f - V_\varphi^g p_k\|_{B_\psi} \leq \|V_\varphi^g\|\|f - p_k\|_{L^2_\psi} \to 0 (k \to \infty).$$

Since $V_\varphi^g p_k \in B_{\psi,0}(\mathbb{D})$ and $B_{\psi,0}(\mathbb{D})$ is the the closed subset of $B_\psi(\mathbb{D})$, we have $V_\varphi^g(AL^2_\psi(\mathbb{D})) \subset B_{\psi,0}(\mathbb{D})$.
Since $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded , we see that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded. □

Theorem 3.3. *Let φ be an analytic self-map of the unit disk $\mathbb{D}$ and $g \in H(\mathbb{D})$, $\psi \in \mathcal{D}$. Then $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact if and only if*

$$\lim_{|\varphi(z)|\to 1} e^{\psi(\varphi(z))-2\psi(z)}\sqrt{\Delta\psi(\varphi(z))}|g'(z)| = 0.$$

Proof. By Lemma 2.2,

$$e^{\psi(\varphi(z))}\sqrt{\Delta\psi(\varphi(z))}\sim\sqrt{K(\varphi(z),\varphi(z))},$$

therefore, we should only show that V_φ^g is compact if and only if

$$\lim_{|\varphi(z)|\to 1}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|=0.$$

First, assume that $\lim_{|\varphi(z)|\to 1}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|=0$, then for any $\varepsilon>0$, there is a positive real number $r_0\in(0,1)$ such that

$$\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|<\varepsilon$$

when $r_0<|\varphi(z)|<1$. Besides, $\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|$ is bounded when $|\varphi(z)|\le r_0$, it is easy to see that

$$\sup_{z\in\mathbb{D}}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|<\infty.$$

By Theorem 3.1, $V_\varphi^g: AL^2_\psi(\mathbb{D})\to B_\psi(\mathbb{D})$ is bounded.

Let $\{f_k\}_{k\in\mathbb{N}}$ be a bounded sequence in $AL^2_\psi(\mathbb{D})$ which uniformly converges to zero on any compact subset of $\mathbb{D}$ as $k\to\infty$. Assume that for any $k\in\mathbb{N}$,$\|f_k\|\le C$, for some positive constant C. Note that

$$\sup_{|\varphi(z)|\le r_0}|g'(z)|e^{-2\psi(z)}\le M$$

for some constant M. It follows that

$$\begin{aligned}
|(V_\varphi^g f_k)'(z)|e^{-2\psi(z)} &= |f_k(\varphi(z))||g'(z)|e^{-2\psi(z)}\\
&\le \sup_{|\varphi(z)|\le r_0}|f_k(\varphi(z))||g'(z)|e^{-2\psi(z)}\\
&\quad+\sup_{|\varphi(z)|> r_0}|f_k(\varphi(z))||g'(z)|e^{-2\psi(z)}\\
&\le M\sup_{|\varphi(z)|\le r_0}|f_k(\varphi(z))|\\
&\quad+\|f_k\|_{L^2_\psi}\sup_{|\varphi(z)|> r_0}\sqrt{K(\varphi(z),\varphi(z))}|g'(z)|e^{-2\psi(z)}\\
&\le M\sup_{|\varphi(z)|\le r_0}|f_k(\varphi(z))|+\varepsilon\|f_k\|_{L^2_\psi}\\
&\le M\sup_{|\varphi(z)|\le r_0}|f_k(\varphi(z))|+\varepsilon C.
\end{aligned}$$

Since $\{f_k\}_{k\in\mathbb{N}}$ uniformly converges to zero on any compact subset of $\mathbb{D}$, there exists a $K\in\mathbb{Z}^+$ such that if $k>K$, we have $\sup_{|\varphi(z)|\le r_0}|f_k(\varphi(z))|<\varepsilon$. Therefore,

$\|V_\varphi^g f_k\|_{B_\psi} \to 0$ as $k \to \infty$. By Lemma 2.4, we see that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact.

Conversely, suppose that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact, then it is clear that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is bounded. Let $\{z_k\}_{k\in\mathbb{N}}$ be sequence in $\mathbb{D}$ such that $\lim\limits_{k\to\infty} |\varphi(z_k)| = 1$. Let

$$f_k(z) = \frac{K(z, \varphi(z_k))}{\sqrt{K(\varphi(z_k), \varphi(z_k))}},$$

then, $f_k \in AL^2_\psi(\mathbb{D})$ and $\|f_k\|_{L^2_\psi} = 1$.

Since the set consisting of polynomials is dense in $AL^2_\psi(\mathbb{D})$, for any $\varepsilon > 0$ and $f \in AL^2_\psi(\mathbb{D})$, there exists a polynomial $P_{f,\varepsilon}(z) \in AL^2_\psi(\mathbb{D})$ such that

$$\|P_{f,\varepsilon} - f\|_{L^2_\psi} < \frac{\varepsilon}{2}.$$

As

$$\begin{aligned} |\langle f_k, f\rangle| &\leq |\langle f_k, f - P_{f,\varepsilon}\rangle| + |\langle P_{f,\varepsilon}, f_k\rangle| \\ &\leq \|f_k\|_{L^2_\psi}\|f - P_{f,\varepsilon}\|_{L^2_\psi} + |\langle P_{f,\varepsilon}, f_k\rangle| \end{aligned}$$

by Lemma 2.2 and Definition 1.1, we have

$$\sqrt{K(z,z)} \geq C_1 \tau(z)^{-1} e^{\psi(z)} \geq \frac{C_2 e^{\psi(z)}}{1 - |z|}$$

for some positive constants C_1 and C_2. Notice that $\lim_{k\to\infty} |\varphi(z_k)| = 0$, we have

$$\langle P_{f,\varepsilon}, f_k\rangle = \langle P_{f,\varepsilon}, \frac{K_{\varphi(z_k)}}{\|K_{\varphi(z_k)}\|_{L^2_\psi}}\rangle = \frac{1}{\|K_{\varphi(z_k)}\|_{L^2_\psi}} P_{f,\varepsilon}(\varphi(z_k)) \to 0 \quad (k \to \infty)$$

That means that f_k weakly converges to zero as $k \to \infty$.

Because $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact, we see that $\|V_\varphi^g f_k\|_{B_\psi} \to 0 \quad (k \to \infty)$. From the following fact

$$\begin{aligned} \|V_\varphi^g f_k\|_{B_\psi} &\geq \sup_{z\in\mathbb{D}} |f_k(\varphi(z))||g'(z)|e^{-2\psi(z)} \\ &\geq |f_k(\varphi(z_k))||g'(z_k)|e^{-2\psi(z_k)} \\ &= \sqrt{K(\varphi(z_k), \varphi(z_k))}|g'(z_k)|e^{-2\psi(z_k)} \end{aligned}$$

we immediately obtain that $\lim\limits_{|\varphi(z)|\to 1} \sqrt{K(\varphi(z), \varphi(z))}|g'(z)|e^{-2\psi(z)} = 0$. The proof is completed. □

Theorem 3.4. *Let φ be an analytic self-map of the unit disk $\mathbb{D}$ and $g \in H(\mathbb{D})$, $\psi \in \mathcal{D}$. Then the operator $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is compact if and only if $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded and*

$$\lim_{|z|\to 1} e^{\psi(\varphi(z))-2\psi(z)}\sqrt{\Delta\psi(\varphi(z))}|g'(z)| = 0.$$

Proof. At first, we note that

$$\lim_{|z|\to 1} e^{\psi(\varphi(z))-2\psi(z)}\sqrt{\Delta\psi(\varphi(z))}|g'(z)| = 0$$

equals to

$$\lim_{|z|\to 1} \sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| = 0.$$

Firstly, we prove the sufficiency . Let $K = \{f : f \in AL^2_\psi(\mathbb{D}), \|f\|_{L^2_\psi} \le 1\}$. As $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded, $\{V_\varphi^g f : f \in K\}$ is the bounded closed set of $B_{\psi,0}(\mathbb{D})$. It suffices to show that $\{V_\varphi^g f : f \in K\}$ is compact in $B_{\psi,0}(\mathbb{D})$. By Lemma 2.5, it is only to prove

$$\lim_{|z|\to 1}\sup_{\|f\|_{L^2_\psi}\le 1} e^{-2\psi(z)}|(V_\varphi^g f)'(z)| = 0.$$

By lemma 2.3, for any $f \in K$, we have

$$\begin{aligned} e^{-2\psi(z)}|(V_\varphi^g f)'(z)| &= e^{-2\psi(z)}|f(\varphi(z))||g'(z)| \\ &\le \|f\|_{L^2_\psi}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| \\ &\le \sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)|. \end{aligned}$$

Note that the condition $\lim_{|z|\to 1}\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| = 0$, we have

$$\lim_{|z|\to 1}\sup_{\|f\|_{L^2_\psi}\le 1} e^{-2\psi(z)}|(V_\varphi^g f)'(z)| = 0.$$

Therefore, the operator $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is compact.

Secondly, we will prove the necessity. Suppose $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is compact, it is obvious that $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_\psi(\mathbb{D})$ is compact. By Theorem 3.3, we have

$$\lim_{|\varphi(z)|\to 1} \sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| = 0.$$

That is, for any $\varepsilon > 0$, there exists an $r \in (0,1)$, such that

$$\sqrt{K(\varphi(z),\varphi(z))}e^{-2\psi(z)}|g'(z)| < \varepsilon. \qquad (*)$$

when $r < \varphi(z) < 1$.

Since $V_\varphi^g : AL^2_\psi(\mathbb{D}) \to B_{\psi,0}(\mathbb{D})$ is bounded, by Theorem 3.2, we have

$$\lim_{|z|\to 1} e^{-2\psi(z)}|g'(z)| = 0.$$

Let $\varepsilon' = \frac{\varepsilon}{C_r}$, there exists a positive real number $\sigma > 0$, such that

$$e^{-2\psi(z)}|g'(z)| < \frac{\varepsilon}{C_r}$$

when $\sigma < |z| < 1$, where C_r is the upper bound of $\sqrt{K(\varphi(z), \varphi(z))}$ when $|\varphi(z)| \leq r$. Therefore, we have

$$\sqrt{K(\varphi(z), \varphi(z))}e^{-2\psi(z)}|g'(z)| < C_r \frac{\varepsilon}{C_r} = \varepsilon.$$

when $\sigma < |z| < 1$ and $|\varphi(z)| \leq r$.

Combines with $(*)$, we see that $\sqrt{K(\varphi(z), \varphi(z))}e^{-2\psi(z)}|g'(z)| < \varepsilon$ when $\sigma < |z| < 1$. Therefore,

$$\lim_{|z|\to 1} \sqrt{K(\varphi(z), \varphi(z))}e^{-2\psi(z)}|g'(z)| = 0.$$

The proof is completed. □

References

[1] Peng Lin and R. Rochberg, Hankel operators on the weighted Bergman spaces with exponential type weights, Integr. Equ. Oper. Teory, 21(1995), 460-483.

[2] Peng Lin and R. Rochberg, Trace ideal Criteria for Toeplitz and Hankel operators on the weighted Bergman spaces with exponential type weights, Pacific Journal of Mathematics, Vol.173(1996), 127-146.

[3] Songxiao Li. Volterra composition operators between weighted Bergman spaces and Bloch type spaces, J. Korean Math. Soc, 45(2008), 229-248.

[4] Jordi Pau, Jose Angel Pelaez. Volterra type operators on Bergman spaces with exponential type weights, Mathematics Subject Classification, (2011), 1-14.